



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

My first impression was utter astonishment at encountering in the eastern United States a bird the appearance of which was so unfamiliar to me. A yellow breast, differentiated from white throat and belly, bold white eye-stripe contrasted with black, rather large slightly curved bill, and other characters having been noted, I was able to immediately identify my bird with certainty in a text-book courteously loaned me by the director of the Miami Aquarium. Its identity was further verified a few days later at Nassau, Bahamas, where I found the same species common enough.

At Nassau, one of these birds was observed taking a morning bath, fluttering in the dew on the broad leaves of a low plant. This is likely a common habit, but was none the less interesting to observe for the first time.—J. T. NICHOLS, *New York City*.

The Sycamore Warbler (*Dendroica dominica albilora*) on the Coast of South Carolina.—I wish to place on record the capture by myself of a young female in full autumnal plumage of this form on the morning of July 13, 1916, near Mount Pleasant. Long before I left home for a swamp (where I collect), a storm was prevailing which increased to hurricane violence before sunset. This bird was carefully examined after being shot and there was the faintest trace of yellow in the supercilary stripe. I, however, waited until I could collect another female of corresponding age to establish the identity to a certainty and found that I had taken the western form of *dominica*. Of the very large series of *D. dominica dominica* that Mr. Brewster collected near Charleston in 1883, '84 and '85, not a single individual showed any tendency towards *albilora*, if my memory serves me correctly, he and I having remarked upon it when the birds were collected.

Mr. Leverett Mills Loomis found the Sycamore Warbler to be a regular fall migrant at Chester, S. C., and noted it as breeding in Pickens County, S. C.

The migration of this subspecies is truly remarkable for the specimen taken by me on July 13, 1916, is the first one I have ever seen since I began to collect birds in 1883.—ARTHUR T. WAYNE, *Mount Pleasant, South Carolina*.

The Orange-crowned Warbler in Indiana.—On the eighth of May, 1921, I took a female Orange-crowned Warbler, *Vermivora celata*, at Dune Park, Porter County, Indiana. This little bird was industriously exploring the terminal clumps of a thicket of low willows in true Kinglet fashion and I came near passing it by as such until attracted by its very yellowish appearance. Captures of the Orange-crowned Warbler in the east Central States are infrequent and I wonder if this fact is not due in part to the bird being overlooked and not entirely to its extreme rarity.—CHRESWELL J. HUNT, *Chicago, Illinois*.